

# New and Emerging Drug Molecules Against Obesity

Journal of Cardiovascular  
Pharmacology and Therapeutics  
2014, Vol 19(1) 65-76  
© The Author(s) 2013  
Reprints and permission:  
sagepub.com/journalsPermissions.nav  
DOI: 10.1177/1074248413501017  
cpt.sagepub.com



Melvin George, MD, DM<sup>1</sup>, Muthukumar Rajaram, MPharm<sup>1</sup>,  
and Elangovan Shanmugam, MD, DM<sup>1</sup>

## Abstract

Obesity has become a growing pandemic of alarming proportions in the developed and developing countries over the last few decades. The most perturbing fact regarding obesity is the increased predisposition for coronary artery disease, congestive heart failure and sudden cardiac death. The modest efficacy of current anti-obesity agents such as orlistat and the increasing withdrawals of several anti-obesity agents such as sibutramine, rimonabant have led to huge gaps in the pharmacotherapy of obesity. Lorcaserin and Phentermine-topiramate combination (phen-top) are two drugs approved by US FDA in 2012. Lorcaserin, a 5HT<sub>2C</sub> agonist has moderate efficacy with an acceptable safety profile. Clinical trials with Phen-top have shown a reasonable efficacy but at the cost of risks such as teratogenicity and psychiatric disturbances. Cetilistat, a lipase inhibitor is claimed to have superior safety profile to orlistat and is in phase 3 clinical trials. Other promising anti-obesity molecules acting on the gut which are in clinical trials include exenatide and liraglutide. Drugs which act on the monoaminergic and opioid systems include bupropion-naltrexone and bupropion-zonisamide. Other novel first-in-class drugs which have been explored and have limited success in early clinical development include velneperit, tesofensine, and beloranib. Tesofensine is a triple monoamine re-uptake inhibitor, velneperit acts as a neuropeptide Y5 receptor antagonist and beloranib is a methionine amino peptidase 2 inhibitor. Novel targets such as histamine H3 receptor, VEGF, matrix-metalloproteinase, sirtuin receptors are also being investigated. This review is an attempt to describe the new and emerging molecules that are in clinical development for obesity.

## Keywords

new drugs, emerging drugs, obesity, weight loss, lorcaserin, phentermine-topiramate

## Introduction

Obesity is a growing pandemic that has risen to alarming proportions over the last decade in the developed world. The situation is also changing in the developing world with the increasing westernization of food habits and lifestyle. According to the World Health Organization projections, by 2015 approximately 2.3 billion adults will be overweight and more than 700 million will be obese.<sup>1</sup> In addition, the economic cost of obesity-associated diseases is approaching US\$100 billion per year in the United States.<sup>2</sup> The most perturbing fact regarding obesity is the increased predisposition for coronary artery disease, congestive heart failure, and sudden cardiac death. Besides cardiovascular (CV) risk, overweight (body mass index [BMI] 25-29) and obesity (BMI >29) are major risk factors for a wide range of diseases such as diabetes, dyslipidemia, obstructive sleep apnea, asthma, musculoskeletal disorders, and cancers.<sup>3</sup> There is compelling evidence to suggest that obesity is associated with more morbidity than smoking, alcoholism, and poverty and is on the verge of overtaking smoking as a leading cause of preventable death in the United States.<sup>4,5</sup>

The current methods for reducing body weight include diet, exercise, drug therapy, bariatric surgery, or combinations of those

mentioned previously.<sup>6</sup> Unfortunately, failure to maintain changes in diet and exercise patterns is a major reason for weight regain experienced by most people. Bariatric surgery, generally used for patients with morbid obesity (BMI  $\geq$  40), is effective for sustaining weight loss but is often fraught with catastrophic complications such as increased mortality.<sup>7,8</sup> The drugs that are currently approved for obesity by the US Food and Drug Administration (FDA) include orlistat, phendimetrazine, and diethylpropion, of which only orlistat is approved for long-term use. Although these drugs have existed in the market for several years, it has failed to make any significant impact on the burden of obesity and its consequences, especially on the CV health. Table 1 highlights the limitations of these older antiobesity agents

<sup>1</sup> Department of Cardiology, SRM Medical College Hospital & Research Centre, Kanchipuram, Tamil Nadu, India

**Manuscript submitted:** June 4, 2013; **accepted:** July 17, 2013.

### Corresponding Author:

Melvin George, Department of Cardiology, SRM Medical College Hospital & Research Centre, Kanchipuram 603203, Tamil Nadu, India.  
Email: melvingeorge2003@gmail.com

**Table I.** Demerits of Older Antiobesity Agents in Clinical Practice.<sup>9-12</sup>

Drug	Dosage	Mechanism of Action	Demerits
Orlistat	120 mg oral 3 times daily	Lipase inhibitor	<ul style="list-style-type: none"> <li>– Serious hepatic effects such as cholelithiasis, cholestatic hepatitis, and subacute liver failure known to occur</li> <li>– FDA warning on liver injury potential</li> </ul>
Diethylpropion	75 mg (SR) once daily, 1 hour before meals	Amphetamine derivative, causes appetite suppression	<ul style="list-style-type: none"> <li>– Safety in pregnant and breast feeding women is not known</li> <li>– Schedule IV controlled substance</li> <li>– Available for short-term use only</li> <li>– Clinical studies negligible</li> </ul>
Phendimetrazine	105 mg (extended release capsule) once daily, 30-60 minutes before morning meal	Amphetamine derivative, causes appetite suppression	<ul style="list-style-type: none"> <li>– Causes adverse effects, including pulmonary arterial hypertension, valvulopathy, and the potential for abuse and dependency</li> <li>– Rhabdomyolysis, interstitial nephritis reported</li> <li>– Available for short-term use only</li> <li>– Schedule IV controlled substance</li> <li>– Clinical studies negligible</li> </ul>

Abbreviations: FDA, Food and Drug Administration; SR, sustained release.

along with their regulatory status.<sup>9-12</sup> Thus, there is an obvious need for an effective antiobesity agent that causes substantial weight loss thereby reducing the risk of cardiovascular disease (CVD). This article is an attempt to highlight the new molecules that have been recently approved for obesity by the US FDA and the emerging molecules that are in clinical development for obesity.

## Antiobesity Molecules Recently Approved by the US FDA

### Lorcaserin

Lorcaserin is a new molecular entity with a 2-ringed structure that has high affinity for the 5-HT<sub>2C</sub> receptor that is present in the hypothalamic proopiomelanocortin neurons. Chemically, lorcaserin is described as [1R]-8-chloro-1-methyl-2,3,4,5-tetrahydro-1H-3-benzazepine hydrochloride.<sup>13</sup> On account of its low-molecular weight, nonpolymer structure, and a binding activity that alters biotarget functions, it is considered a “small molecule.” Binding of lorcaserin to the 5-HT<sub>2C</sub> receptors in proopiomelanocortin neurons promotes satiety and reduces the individual’s food intake. Lorcaserin is selective for the 5-HT<sub>2C</sub> subtype and does not have significant activity at either 5-HT<sub>2B</sub> or 5-HT<sub>2A</sub> receptors; activity at the 5-HT<sub>2B</sub> receptor has been associated with the development of valvular heart disease with earlier antiobesity agents such as dexfenfluramine.<sup>14,15</sup>

**Pharmacokinetics of Lorcaserin.** The recommended clinical dose of lorcaserin is 10 mg twice daily. Following oral administration, more than 90% of lorcaserin is rapidly absorbed. The time taken to attain maximum plasma concentration is approximately 1.5 to 2 hours, and its half-life ( $t_{1/2}$ ) is about 11 hours. Approximately 70% of the drugs are bound to plasma proteins. The volume of distribution in the body is ~252 L in a 92.5-kg patient. Lorcaserin is mainly metabolized in the liver and excreted by the kidney. The major circulating metabolite is the

sulfamate of lorcaserin (M1), and the major urinary metabolite is N-carbamoyl glucuronide (M5).<sup>13-17</sup>

**Efficacy of Lorcaserin.** Lorcaserin has been evaluated in 3 large randomized, placebo-controlled, double-blind studies, the results of which paved way for the drug to receive an FDA approval in June 27, 2012, for the treatment of obesity. The first study investigating lorcaserin, the Behavioral Modification and Lorcaserin for Overweight and Obesity Management (BLOOM) study, enrolled people 18 to 65 years of age with a baseline BMI of 30 to 45 kg/m<sup>2</sup> or a BMI of 27 to 45 kg/m<sup>2</sup> with at least 1 concomitant weight-related comorbidity, such as hypertension, CVD, dyslipidemia, impaired glucose tolerance, or obstructive sleep apnea.<sup>18</sup> A total of 3182 participants were randomized to receive lorcaserin 10 mg twice a day (n = 1595) or placebo (n = 1587) for 52 weeks. At 52 weeks, those receiving placebo continued for an additional 52 weeks, whereas those receiving lorcaserin twice a day were randomized in a 2:1 ratio to continue receiving lorcaserin or be switched to placebo. The primary outcome, which was the percentage of patients who achieved 5% weight loss of baseline weight after 52 weeks, occurred in 47.5% of those receiving lorcaserin compared with 20.3% receiving placebo (modified intention to treat [mITT] analysis population with last observation carried forward [LOCF],  $P < .001$ ). The mean decrease in weight (calculated using the least squares mean) was 5.8% (5.8 kg) for lorcaserin compared with 2.2% (2.2 kg) for placebo. After 104 weeks of treatment, more patients who continued taking lorcaserin were able to maintain weight loss of 5% compared with those who transitioned to placebo (67.9% vs 50.3%,  $P < .001$ ). Secondary outcomes, which included percentage achieving >10% weight loss ( $P < .001$ ), reduction in waist circumference, blood pressure reduction, and BMI, were improved with lorcaserin after 52 weeks of treatment, although the clinical significance of these differences is not known.

In the BLOSSOM trial, a total of 4008 participants were randomized to receive either lorcaserin 10 mg (n = 801),

lorcaserin 10 mg twice a day ( $n = 1602$ ), or placebo ( $n = 1601$ ) for 52 weeks in addition to nutritional and exercise counseling provided at each study visit.<sup>19</sup> After 1 year of treatment, 47.2% of those receiving lorcaserin twice a day lost >5% of their baseline weight compared with 40.2% of those receiving lorcaserin every day and 25% of those receiving placebo ( $P < .001$  for both doses of lorcaserin vs placebo). Greater than 10% weight loss was achieved by 22.6% receiving lorcaserin twice a day, 17.4% receiving lorcaserin every day, and 9.7% receiving placebo ( $P < .001$  for both doses vs placebo). Absolute weight loss, calculated by least squares mean, was 5.8, 4.7, and 2.9 kg for lorcaserin twice a day, lorcaserin every day, and placebo, respectively. Although systolic blood pressure, diastolic blood pressure, and heart rate decreased in all the groups, the difference was not statistically significant. Importantly, no differences were noted between lorcaserin and placebo in echocardiographic outcomes.

The BLOOM in Diabetes Mellitus (BLOOM-DM) study enrolled people aged 18 to 65 years with a BMI of 27 to 45 kg/m<sup>2</sup> and a diagnosis of type 2 diabetes with a hemoglobin A1c (HbA1c) level of 7% to 10%.<sup>20</sup> Patients were stratified based on baseline type 2 diabetes treatment and randomized to lorcaserin 10 mg twice a day ( $n = 256$ ), lorcaserin 10 mg every day ( $n = 95$ ), or placebo ( $n = 253$ ) for 52 weeks; enrollment into the lorcaserin 10 mg every day group was stopped after 8 months because of slow enrollment. Patients received standardized nutritional and exercise counseling at each visit. At baseline, the mean body weight was 104 kg, and the mean BMI was 36 kg/m<sup>2</sup>. Approximately 45% of the patients in the lorcaserin once daily group and 38% of the lorcaserin twice daily group achieved at least 5% weight loss compared with only 16% in the placebo group ( $P < .001$  for both doses compared with placebo). A 10% reduction in weight was achieved by only 4.4% in the placebo group compared with 16.3% of those receiving lorcaserin twice daily and 18.1% of those receiving lorcaserin once daily. ( $P < .001$  for both doses compared with placebo). Absolute weight loss (using least squares means) was 4.7 kg for lorcaserin twice daily, 5 kg for lorcaserin once daily, and 1.6 kg for placebo. Incidence of FDA-defined valvular disease did not differ between the groups.

In February 2007, a draft guidance document was released by the US FDA on the principles of developing drugs for weight loss management. In order for an antiobesity agent to be deemed effective, it needs to fulfill certain benchmarks as laid down by the guidelines. Treatment with the antiobesity molecule for 1 year should result in at least a 5% difference between the active group and the placebo, and this difference must be statistically significant or the proportion of patients who lose  $\geq 5\%$  of baseline body weight in the active product group should be at least 35%, and should approximately double the proportion in the placebo-treated group, and the difference between the groups should be statistically significant. By these standards, the efficacy of lorcaserin that was achieved in the three phase 3 clinical trials was only modest. Additionally, the high percentage of dropouts in all the 3 studies may actually tilt the results in favor of the drug, since the statistical

approach used was mITT with LOCF. It is not known whether the results would appear different, if an imputation analysis was performed instead of the former. Nevertheless, considering the fact that the currently available antiobesity agents have only a similar or slightly lower efficacy, the drug regulators in the US FDA have approved the drug for marketing.

**Safety of Lorcaserin.** Lorcaserin had to face several regulatory hiccups prior to its approval by FDA on account of several safety concerns. Initially, FDA rejected an application for approval of this drug due to concerns of unexplained preclinical carcinogenicity signals in the form of breast tumors in rats treated with lorcaserin. Another disturbing feature with lorcaserin was the rising rates of new valvulopathy. However, subsequent analysis has shown that the absolute numbers of valvulopathy were much less. The most common adverse events with lorcaserin include headache, nausea, dizziness, dry mouth, paresthesia, upper respiratory tract infection, and anorexia. A small number of patients did experience a change in mood.<sup>21</sup> There have not been any carcinogenic signals from the data generated from human studies as of date.

On the basis of the results of these studies, lorcaserin is approved at a dose of 10 mg twice daily in patients with a BMI of 30 or 27 kg/m<sup>2</sup> with at least 1 weight-related comorbidity, such as hypertension, type 2 diabetes, or dyslipidemia, in addition to a reduced calorie diet (RCD) and increased physical activity. The labeling indicates response to therapy should be assessed at week 12, and if there is a 5% decrease in weight, use of the drug should be discontinued, because it will be unlikely that the patient will achieve and sustain adequate weight loss with continued treatment. The long-term CV safety of lorcaserin with respect to major adverse cardiac events such as stroke and myocardial infarction (MI) is currently being investigated in postmarketing studies.<sup>22</sup> Although the drug has only a modest efficacy, its favorable safety profile may make it a preferred choice in obesity clinics in the years ahead.

**Phentermine–Topiramate.** The phentermine–topiramate (Phen-top) combination is a once-daily formulation designed to provide an immediate release of phentermine and a delayed release of topiramate. Phentermine reduces the food intake by enhancement of norepinephrine release and blockade of norepinephrine reuptake. The antiobesity action of topiramate although largely unexplained is believed to be due to increased energy expenditure, decreased energetic efficiency, and decreased caloric intake from preclinical studies. The combination allows lower doses of the individual compounds and is reported to cause fewer side effects than the individual compounds. The fixed dose combination was approved by the FDA in July 17, 2012, for promoting weight loss based on the tolerability and efficacy data from 3 large randomized, double-blind, placebo-controlled trials.

The EQUIP trial (Controlled-Release Phentermine/Topiramate in Severely Obese Adults: A Randomized Controlled Trial) included people aged 18 to 70 years with a BMI of 35 kg/m<sup>2</sup> with no upper limit, a fasting blood glucose level 100

**Table 2.** Newly Approved Antiobesity Agents.

Drug	Dosage	Mechanism of Action	Common Adverse Events
Lorcaserin	10 mg oral twice daily	5-HT <sub>2c</sub> agonist	Headache, upper respiratory tract symptoms and infection, dizziness, nausea, constipation, and fatigue
Phentermine–topiramate	7.5 mg of phentermine and 46 mg of ER topiramate once daily	Amphetamine derivative	Dry mouth, unpleasant taste, diarrhea, constipation, vomiting, headache, insomnia, shortness of breath, and chest pain

Abbreviations: ER, extended release.

mg/dL, and a blood pressure 140/90 mm Hg.<sup>23</sup> Participants were stratified by sex and randomized to controlled-release phen-top (phen-top CR) 15/92 mg (n = 512), phen-top CR 3.75/23 mg (n = 241), or placebo (n = 514) for 52 weeks. The primary outcome was percentage of weight loss at week 52; analysis was by ITT with LOCF. The percentage of weight loss with high-dose phen-top was 10.9% compared with 5.1% in the low-dose phen-top and 1.6% for placebo ( $P < .001$  for both doses compared with placebo). Absolute weight loss based on least squares mean was 12.6 kg for high-dose phen-top, 6 kg for low-dose phen-top CR, and 1.8 kg for placebo.

The CONQUER study (Effects of Low-Dose, Controlled-Release, Phentermine Plus Topiramate Combination on Weight and Associated Comorbidities in Overweight and Obese Adults) included people with a BMI of 27 to 45 kg/m<sup>2</sup> or those with type 2 diabetes with any BMI.<sup>24</sup> Participants were stratified based on the gender and presence of type 2 diabetes and were randomized to phen-top CR 7.5/46 mg (n = 498), phen-top CR 15/92 mg (n = 995), or placebo (n = 994) which was continued for 52 weeks. The coprimary outcome of the CONQUER study was mean percentage change in weight and percentage of patients achieving at least 5% weight loss; analysis was by ITT with LOCF. The percentage weight loss was 7.8% (8.1 kg) with low-dose phen-top, 9.8% (10.2 kg) for high-dose phen-top, and 1.2% (1.8 kg) for placebo ( $P < .001$  for both doses vs placebo). Approximately 21%, 62%, and 70% achieved at least 5% weight loss while receiving placebo, low-dose phen-top, and high-dose phen-top CR, respectively ( $P < .001$  for both doses vs placebo). Treatment with phen-top CR improved blood pressure, high-density lipoprotein cholesterol, triglyceride (TG), HbA1c, and fasting glucose compared with placebo ( $P < .05$  for all comparisons), although the absolute differences are of questionable clinical significance.

The SEQUEL study was a 1-year extension of the CONQUER trial; patients were only eligible if they completed the CONQUER study on treatment and complied with protocol requirements. Patients continued their randomized treatment from CONQUER for an additional 52 weeks. In the SEQUEL study, the high-dose group achieved a weight loss percentage of 10.5% (10.9 kg), and this benefit was also observed in the low-dose group [9.3% (9.6 kg)] compared to placebo [1.8% (2.1 kg);  $P < .001$  for both doses vs placebo]. Nearly 80% of participants in the high-dose group attained 5% weight loss compared to 75% in the low-dose group and 30% receiving

placebo ( $P < .001$  for both doses compared with placebo). The reduction was sustained even after patients continued the treatment for the second consecutive year. However, the study investigators do acknowledge the inherent bias of the SEQUEL study conclusions since only the adherent patients of the CONQUER study were chosen to participate, thus representing a group that is already well satisfied with therapy.<sup>25</sup>

**Safety.** Although the study investigators were able to document the efficacy of phen-top combination, there were many questions raised by the FDA regarding the safety of the molecule.<sup>20</sup> The most frequently reported adverse events associated with phen-top treatment were paresthesia (17.0%), dry mouth (16.6%), constipation (15.1%), upper respiratory tract infection (13.5%), nasopharyngitis (10.0%), and headache (9.8%). There were concerns of teratogenic risk in women with child-bearing potential receiving topiramate. A retrospective observational study named Fetal Outcome Retrospective TopiRamate Exposure Study (FORTRESS) was initiated by the manufacturing agency, using electronic medical health records, to determine the fetal outcome in the offspring of mothers exposed to topiramate. The data showed that exposed women had nearly a 2 times greater risk of having children with oral clefts.<sup>26</sup> There was also a concern about the increase in heart rate with phen-top and whether this increase could lead to increased CV risk, as seen with sibutramine. This concern was laid to rest when data analysis revealed that there were only a small number of major adverse cardiac events among patients exposed to phen-top, and these patients did not have increased heart rates. Taking these factors into consideration and the beneficial effect of phen-top on the BMI, blood pressure, and diabetic status, the CV safety of phen-top appears reasonable. Nevertheless, there is an onus on the regulators to seek data from prospective long-term studies to assess the effects of the drug on the risk of MI, stroke, CV, and all-cause mortality.

On the basis of the results of these studies, phen-top CR was approved by the US FDA for the treatment of obesity in patients with a BMI of 30 or 27 kg/m<sup>2</sup> with at least one weight-related comorbidity, such as hypertension, type 2 diabetes mellitus, or dyslipidemia, in addition to an RCD and increased physical activity. The Phen-top CR therapy should be started at 3.75/23 mg, taken once daily in the morning, and after 14 days the dose can be increased to 7.5/46 mg. If after 12 weeks, at least 3% weight loss has not been achieved, use of the

drug can be discontinued or the dose can be increased further to 11.25/69 mg for 14 days followed by a final dose increase up to 15/92 mg. Weight loss should be evaluated after an additional 12 weeks, and if 5% weight loss is not achieved, therapy should be discontinued. The Risk Evaluation and Mitigation Strategy has been put in place by the US FDA for phen-top to ensure that the benefits of the drug overcome the risks associated with it such as teratogenicity. Women of child-bearing potential are cautioned regarding the risks such as orofacial clefts to the developing fetus with the use of the drug, counseled on the use of effective contraception during drug use, and to terminate the drug if pregnancy does occur.<sup>27</sup> The drug is yet to receive European Medicines Agency approval. Table 2 summarizes the salient features of lorcaserin and phen-top.

## Antiobesity Molecules in Clinical Development

### Agents Acting Through Central Mechanisms

**Bupropion–Naltrexone.** Bupropion is an atypical anti-depressant that inhibits the reuptake of dopamine and noradrenaline. Although the drug was detected to have weight loss-inducing potential, the decrease in body weight was a mere 2.8 kg after 6 months of therapy, and hence the drug did not match the FDA standards as monotherapy for obesity. Bupropion additionally stimulates proopiomelanocortin firing in the arcuate nucleus of the hypothalamus which in turn releases  $\alpha$ -melanocyte-stimulating hormone that mediates the anorectic effects. However, the stimulation of POMC also stimulates a negative-feedback loop that releases  $\beta$  endorphins, which suppress further POMC firing. One of the reasons attributed to the failure of bupropion as a sole agent for obesity is the negative-feedback loop activation by bupropion. This negative feedback can be suppressed by naltrexone, a  $\mu$ -opioid receptor antagonist.

A randomized, placebo-controlled trial by Wadden et al in 793 patients demonstrated that weight loss was  $5.1\% \pm 0.6\%$  with placebo + behavioral modification (BM) versus  $9.3\% \pm 0.4\%$  with naltrexone–bupropion combination + BM ( $P < .001$ ).<sup>28</sup> In a phase 3 clinical trial carried out in 1742 patients by Greenway et al, patients were randomized to receive naltrexone 32 mg + bupropion; naltrexone 16 mg + bupropion; or placebo. The mean change in body weight was  $-6.1\%$ ,  $-5.0\%$ , and  $-1.3\%$  in the 3 groups, respectively ( $P < .001$  in both groups compared to placebo).<sup>29</sup> The common adverse reactions that were seen among the trial participants receiving the combination therapy included nausea, headache, constipation, dizziness, vomiting, and dry mouth. Another phase 3 clinical trial performed in 1496 patients by Apovian et al showed that patients on NB-32 combination had a weight loss superior to placebo ( $-6.5\%$  vs  $-1.9\%$ ;  $P < .001$ ). More NB-32-treated participants ( $P < .001$ ) experienced  $\geq 5\%$  weight loss versus placebo at week 28 (55.6% vs 17.5%) and week 56 (50.5% vs 17.1%). There was also an improvement in various cardiometabolic risk markers, participant-reported, weight-related quality of life, and control of eating with NB-32.<sup>30</sup> The drug was in the verge of obtaining

FDA approval. However, the regulatory body insisted that the manufacturer carry out long-term CV safety studies before approval.<sup>31</sup>

**Zonisamide–Bupropion.** Zonisamide is an antiepileptic agent that is used in the management of partial seizures. The serendipitous discovery of zonisamide's weight loss promoting potential in patients receiving the drug led the manufacturers to develop it for the treatment of obesity. Although the precise mechanism is yet to be elucidated, modulation of sodium channel, carbonic anhydrase inhibition, and enhancement of dopamine and serotonin transmission are said to play a role in inducing weight loss. Bupropion causes weight loss by increase in the levels of dopamine that decreases appetite. The detrimental effects of zonisamide such as depression and sedation may be overcome by its combination with bupropion. Conversely, the seizure-inducing property of bupropion could be offset by zonisamide. Combination of zonisamide with bupropion was observed to be superior to bupropion or zonisamide monotherapy in inducing weight loss in a pilot study performed among 18 obese women.<sup>32</sup> A phase 2b study clinical trial in 729 patients with obesity showed that the combination in 2 different doses had a weight loss that was superior to placebo (6.1% and 7.5% vs 1.4%,  $P < .01$ ).<sup>33,34</sup> Nausea, headache, and insomnia were the main adverse events in this clinical trial with the combination. The manufacturer Orexigen is in the process of initiating phase 3 clinical trials with the fixed dose combination.

**Tesofensine.** Tesofensine is a monoamine reuptake inhibitor originally developed for Alzheimer disease and Parkinson disease, which was incidentally found to induce weight loss in clinical trials. The drug prevents the reuptake of serotonin, noradrenaline, and dopamine and thereby suppresses appetite and increases thermogenesis. In a phase 2 clinical trial in which patients were randomized to receive tesofensine or placebo for 24 weeks, tesofensine at a dose of 0.25, 0.5, and 1 mg showed a weight loss of 4.7, 9.1, and 10.6 kg, respectively.<sup>35</sup> The drug was fairly tolerated, with dry mouth, dizziness, constipation, abdominal pain, and nausea being the common side effects. A modest increase in heart rate up to 8 bpm has been recorded with the higher dosage of tesofensine (1 mg).<sup>36</sup> This clinical trial that was published in *Lancet* came under the scrutiny of the Danish drug regulatory agency in 2011, which cited lapses in trial methodologic procedures during a site inspection.<sup>37</sup> The drug is yet to enter confirmatory phase of clinical trial development.

**Velneperit.** It has been well recognized over the last 2 decades that neuropeptide Y (NPY) stimulates food intake, reduces energy expenditure, and increases body weight by activating NPY receptors, Y1, and Y5 present in the hypothalamus. Velneperit is a Y5 receptor antagonist that prevents the binding of NPY to the Y5 receptors and thus decreases hunger and controls energy balance. The RCD study that was carried out in 656 patients randomized them to receive velneperit 800/1600 mg or placebo along with RCD regimen. Patients receiving 800 mg lost an average of 3.8 kg compared to the placebo

group with a mere 0.8 kg ( $P < .0001$ ). In the 800-mg group, 35% of the patients lost greater than 5% of body weight, while in the placebo group only 12% showed >5% reduction. The low-calorie diet (LCD) study which also evaluated velniperit at a dose of 1600 mg showed a weight loss of 7.1 kg in the study group over placebo (4.3 kg). Among the study patients, 52% in the velniperit group and 35% in the placebo group had a weight loss greater than 5%.<sup>33</sup> Nasopharyngitis, sinusitis, upper respiratory infection, sinusitis, and headache were the common adverse events observed. The lack of significant benefit of velniperit over placebo in the LCD study has raised several questions on the future prospects of the drug and the need for further drug development.

**GSK1521498.** Opioid neurotransmission plays a stellar role in behaviors reinforced by primary rewards, especially fatty or sugary foods.<sup>38</sup> Opioid antagonists have been shown to reduce short-term food intake and affective or subjective pleasantness of palatable foods in healthy participants.<sup>39</sup> GSK1521498 is a  $\mu$ -opioid receptor inverse agonist that is being investigated primarily for the treatment of overeating behavior in obesity. A study in healthy volunteers showed that GSK1521498 selectively reduced sensory hedonic ratings of high-sugar and high-fat dairy products and caloric intake of high-fat/high-sucrose snack foods.<sup>40</sup> Another phase 1 study showed the drug to have an acceptable safety profile and pharmacokinetics when the drug was administered for 10 days in multiple doses.<sup>41</sup> It remains to be seen whether the decreased food craving would translate into significant weight reduction when the drug is given for extended periods of time.

### Agents Acting Through Peripheral Mechanisms in the Gut

**Cetilistat.** Cetilistat is a pancreatic lipase inhibitor that is undergoing phase III clinical trials. Pancreatic lipase is a key enzyme that causes breakdown of TGs into free fatty acids that are absorbed in the intestine. Inhibition of pancreatic lipase by cetilistat reduces the conversion of TGs into free fatty acids, and thus the TGs are excreted unchanged in the urine. Cetilistat was administered in 3 different doses of 60, 120, and 240 mg twice daily and compared with placebo. The weight loss obtained was 3.3, 3.5, and 4.1 kg, respectively, at the end of 12 weeks. All 3 groups had a weight loss that was almost similar and was statistically superior to the placebo group.<sup>42</sup> Although cetilistat is similar to orlistat in its mechanism, it is claimed to have a superior safety profile that was demonstrated in a phase 2b clinical trial.<sup>42,43</sup>

**Liraglutide.** In the last decade, glucagon-like peptide 1 (GLP-1) analogs have become a frontline option in the treatment of patients with diabetes. Liraglutide has been the latest of the GLP analogs approved for lowering of blood glucose in diabetes. The GLP-1 enhances insulin secretion, delays gastric emptying, and suppresses appetite. This results in weight loss and decreased energy intake.<sup>44</sup> In a study that evaluated liraglutide in 4 different dosages of 1.2, 1.8, 2.4, or 3 mg, the mean weight loss

obtained in each group at the end of the study was 4.8, 5.5, 6.3, and 7.2 kg, respectively. In the same study, patients in the placebo arm showed a weight loss of 2.8 kg, while those on standard therapy orlistat 120 mg twice daily had a weight loss of 4.1 kg.<sup>45</sup> Liraglutide also showed an acceptable safety profile with the most common adverse reactions experienced being nausea and vomiting. Several clinical trials are currently underway (phases 2 and 3) to test the safety and efficacy of liraglutide as an antiobesity molecule.<sup>46</sup> A meta-analysis showed that treatment with GLP-1 R agonists reduces body weight in both diabetic and nondiabetic patients with obesity.<sup>47</sup> The main limitation with both liraglutide and exenatide is the need for parenteral formulation. Since GLP analogs are currently being explored for their CV risk-lowering potential, this may be an added advantage apart from the weight loss sustained due to drug therapy.

**PF-04620110.** Alteration in fatty acid and TG metabolism is one of the factors implicated in the pathophysiology of obesity. The final step in triacylglycerol formation is catalyzed by the enzyme diacylglycerol acyltransferase 1 (DGAT1), a member of the large family of membrane-bound O-acyltransferases. The DGAT1 causes reesterification of dietary TG in the intestine. PF-04620110 is a selective pharmacological inhibitor of DGAT1. In C57/BL6J mice, acute DGAT1 inhibition alters the temporal and spatial pattern of dietary lipid absorption. The DGAT1 inhibition causes an enrichment of polyunsaturated fatty acids within the TG class of lipids.<sup>48</sup> Phase 1 clinical trials of PF-04620110 have been conducted in healthy volunteers, patients with obesity, and diabetic patients, but the results of these trials could not be retrieved.<sup>49</sup> It is too early to conclude that DGAT inhibition could lead to significant weight reduction; nevertheless, DGAT appears to be evoke interest as an antiobesity and antidiabetic molecule.

### Agents Acting Through Metabolic Pathways in Liver and Adipose Tissue

**Beloranib.** It is a methionine aminopeptidase 2 (MetAP2) inhibitor that has been found to be effective in preclinical studies. The drug inhibits MetAP2 that reduces production of new fatty acid molecules by the liver and helps to convert stored fats into useful energy.<sup>50</sup> A study comparing 3 different intravenous doses of beloranib given twice weekly showed that the median weight reduction was  $-3.8$  kg with beloranib 0.9 mg/meter square versus  $-0.6$  kg with placebo.<sup>51</sup> The drug is currently undergoing phase 2 clinical trials to establish the safety, efficacy, and pharmacokinetics of the drug in patients with obesity.<sup>52</sup> One of the limitations with this molecule is the need for parenteral formulation—subcutaneous formulation and intravenous formulation are presently being studied.

**Resveratrol.** Sirtuin 1 (SIRT1) is an NAD-dependent deacetylase that is upregulated in the setting of calorie restriction in rodents. The SIRT1 binds to PPAR- $\delta$  and suppresses the expression of PPAR- $\delta$  genes that are responsible for fat storage.

**Table 3.** Antiobesity Molecules in Clinical Development.

Drug	Mechanism of Action	Clinical Development Phase	Reference
Cetilistat	Pancreatic lipase inhibitor	NDA filed in Japan	61
Contrave (bupropion/ naltrexone)	Dopamine and noradrenaline reuptake inhibitor/ $\mu$ -opioid receptor agonist	Phase III ongoing	62
Empatic (zonisamide/bupropion)	CA inhibitor/dopamine and noradrenaline reuptake inhibitor	Phase II completed	63
Liraglutide	Long-acting GLP-1 analog	Phase III ongoing	64
Velneperit	Neuropeptide Y5 receptor antagonist	Phase II completed	65
Tesofensine	Triple monoamine reuptake inhibitor	Phase II completed	66
JNJ-16269110	MTP inhibitor	Phase II completed	67
TTP-435	AgRP Inhibitors	Phase II completed	68
Beloranib	MetAP-2 inhibitor	Phase II ongoing	52
Oxyntomodulin	Dual agonist of GLPIR and GCGR	Phase II ongoing	69
GSK1521498	$\mu$ -Opioid inverse agonist	Phase I completed	70
GSK598809	D <sub>3</sub> (dopamine) antagonist	Phase I completed	71
PF-04971729	Selective inhibitor of the sodium-dependent glucose cotransporter 2	Phase I completed	72
PF-04620110	Selective DGAT-1 inhibitor	Phase I completed	73

Abbreviations: AgRP, agouti-related protein; CA, carbonic anhydrase, DGAT-1, diglyceride acyl transferase 1, GLP-1, glucagon-like peptide 1; GCGR, glucagon receptor; MetAP2, methionine aminopeptidase 2; MTP, microsomal triglyceride transfer protein; NDA, new drug application.

It also activates oxidative metabolism and fatty acid oxidation by virtue of its interaction with peroxisome proliferator-activated receptor  $\alpha$  coactivator (PGC-1 $\alpha$ ).<sup>44</sup> Resveratrol is an allosteric activator of SIRT1 that has been demonstrated to protect against insulin resistance and increase the mitochondrial activity in brown adipose tissue and skeletal muscle from preclinical studies. In a double-blind crossover study done in 11 patients with obesity, 30 days of resveratrol supplementation induces metabolic changes such as activation of adenosine monophosphate-activated protein kinase, increased SIRT1 and PGC-1 $\alpha$  protein levels, increased citrate synthase activity without change in mitochondrial content, and improved muscle mitochondrial respiration on a fatty acid-derived substrate.<sup>53</sup> Besides resveratrol, several other sirtuin activators are in early clinical development for obesity.

**ALS-L1023.** The adipose tissue is a highly vascularized organ that is subject to extensive modification in obesity. The growth of adipose tissue is remarkably affected by adipogenesis, angiogenesis, and extracellular matrix (ECM) remodeling. Modulation of ECM also enhances the maturation of microvessels, and this is regulated by matrix metalloproteinase (MMP).<sup>54-56</sup> Active tissue remodeling is associated with adipogenesis and increased MMP expression. The serum concentrations of growth factors such as vascular endothelial growth factor (VEGF), fibroblast growth factor, soluble VEGF receptor 2 (VEGFR2), hepatocyte growth factor, angiopoietin-2, and angiogenin are higher in overweight or obese individuals than in normal-weight individuals.<sup>57</sup> ALS-L1023 is a drug obtained from *Melissa officinalis* that has exhibited enhanced antiangiogenic and MMP inhibitory activities. The messenger RNA-2 expression of angiogenic factors, MMPs, and the resulting adipose tissue growth were modulated by ALS in obese mice.<sup>58</sup> As per the manufacturer Angiolabs, ALS-L-1023 has already reached phase 3 clinical trial

development in Korea.<sup>59</sup> The Ob-X is a mixture of 3 herbs *M officinalis* L (Labiatae), *Morus alba* L (Moraceae), and *Artemisia capillaris* Thunb (Compositae). Administration of Ob-X to ob/ob mice for 5 weeks produced a significant reduction in body weight gain by 27% compared with control. The Ob-X also significantly decreased visceral adipose tissue mass by 15%.<sup>60</sup> The Ob-X is currently being promoted in Korea as a dietary supplement. Thus, it appears that inhibition of MMP and angiogenesis could be a potential target for obesity. Table 3 is an overview of the drugs that are in the pipeline for the treatment of obesity.

**Obesity Paradox.** Although obesity is a strong and well-known risk factor for CVD, there have been some intriguing findings from different investigators that demonstrate the contrary that has been termed obesity paradox. For instance, a study by Uretsky et al in 22 576 treated patients with hypertension showed that all-cause mortality was 30% lower in overweight and patients with obesity, despite the inadequate control of their blood pressure.<sup>83</sup> These findings were replicated in another study performed in 3975 patients, which showed a U-shaped relationship between all-cause, CV and non-CV mortality, and BMI,<sup>84</sup> thereby leading to the hypothesis that obese patients with hypertension have a paradoxically better prognosis. The obesity paradox was also demonstrated in studies that were performed in patients with heart failure.<sup>85</sup> A meta-analysis by Flegal et al also showed that overweight was associated with lower all-cause mortality, while grades 2 and 3 obesity was associated with higher all-cause mortality.<sup>86</sup> This paradoxical relationship is also observed in patients with heart failure having obesity, where higher BMI has been linked with lower mortality. The risk of mortality was lower for every 5-unit increase in BMI ( $P < .001$ ).<sup>87</sup> It is not known whether weight loss-lowering strategies with the new and emerging agents would cause a difference to this obesity paradox. Nevertheless, there is ample evidence

**Table 4.** Antiobesity Drugs That Have Been Withdrawn From the Market.

Drug	Mechanism of Action	Regulatory Status	References
Dinitrophenol	Thermogenesis (uncouples oxidative phosphorylation)	Withdrawn in 1938 due to the risk of neuropathy and cataracts	74, 75, 11
Aminorex	5-HT releaser and reuptake inhibitor; also potent monoamine oxidase inhibitor	Withdrawn in 1968 due to the concern of pulmonary arterial hypertension	76
Fenfluramine and dexfenfluramine	5-HT <sub>2B</sub> receptor agonist	Withdrawn after reports of valvular heart damage and primary pulmonary hypertension in 1997	77-78
Mazindol	Norepinephrine reuptake inhibitor	Withdrawn in 2000 due to an unfavorable risk to benefits ratio	80
Methamphetamine	Appetite suppression	Withdrawn in 2000 due to dependency and abuse potential, cardiovascular adverse effects	76
Phenylpropanolamine	Norepinephrine/dopamine releasing stimulator	Withdrawn for increased risk of hemorrhagic stroke in 2000	81
Phendimetrazine	Appetite suppressant	Currently indicated for the management of exogenous obesity as short-term adjunct, but their use is restricted because of their adverse effects such as pulmonary arterial hypertension, valvulopathy, and the potential for abuse and dependency so it had been withdrawn by European union at the year 2000. Still available in United States	11
Diethylpropion	Appetite suppression	Currently approved drug by US FDA for short-term weight management. But it had been withdrawn by European union at the year 2000 itself due to an unfavorable risk to benefits ratio	80,82
Rimonabant	Selective CBI receptor blocker	Withdrawn in 2009 due to potential of serious psychiatric disorders including depressed mood disorders, anxiety, and suicidal ideation	89,90
Sibutramine	Selective combined serotonin and noradrenaline reuptake inhibitor (appetite suppression)	Withdrawn in 2010 due to increased risk of heart attack and stroke in high-risk patients with cardiac disease	91

Abbreviation: FDA, Food and Drug Administration.

**Table 5.** Antiobesity Agents That Have Failed During Clinical Development.

Drug	Mechanism	Reason for Failure	References
Pramlintide/metreleptin	Amylin analogue/leptin analogue	Commercial reasons	92
Obinipitide	PYY3-36 and pancreatic polypeptide analog	Unknown	93
LY377604	$\beta_3$ adrenergic receptor agonist	Clinical trial terminated due to results from recent nonclinical studies	94
Beta histine	Histamine receptor H <sub>1</sub> agonist and H <sub>3</sub> antagonist	Inconsistent results	95,96
Taranabant	Cannabinoid receptor inverse agonist	Psychiatric side effects	97
MK-0493	Melanocortin receptor 4	Limited efficacy	98

linking weight loss strategies with proven CV benefits.<sup>88</sup> There is a definite need to explore these issues further, and at the moment with our current level of understanding, the most reasonable strategy is to continue achieving weight reduction through pharmacologic and nonpharmacologic therapy.

**Challenges in Obesity Drug Development.** There is almost no other area in drug development, which is replete with as many failures and withdrawals as seen in obesity (Tables 4 and 5).

Some of the major challenges in obesity drug development include the regulatory body's expectations on the risk benefit profile being far superior to other drug classes, considering the high number of drug withdrawals in the past. The lack of widespread reimbursement options for an antiobesity molecule is another serious possible reasons for manufacturers to stay away from antiobesity drug development.<sup>99</sup> The reluctance by many physicians to consider obesity as a disease but more as a risk factor may make them even more hesitant to prescribe antiobesity drug therapy. The regulations also stipulate that antiobesity

agents apart from causing significant weight reduction should also cause improvement in blood pressure, lipids, glycemia, or other beneficial outcomes that are commensurate with the degree of weight loss.<sup>100,101</sup>

## Conclusion

In spite of the checkered past of obesity drug development, with so many promising molecules ending in failure, the search is still on for the elusive molecule that can target obesity and reduce its consequences substantially. Although the regulatory agencies have rightly expressed concern on certain antiobesity molecules and refused marketing approval in lieu of safety issues, this has not blunted the enthusiasm of pharmaceutical companies to discover new molecules, since the market for an antiobesity drug appears too alluring to resist. Lorcaserin and phen-top are the 2 drugs that have been recently approved for obesity. Although not compared from head-to-head clinical trials, phentermine appears to have greater efficacy than lorcaserin. Lorcaserin may be a safer option, more so in women of child-bearing potential due to the teratogenic risk attributed to topiramate. The need of the hour is to gather data from prospective clinical trials that evaluate the long-term CV outcomes with the use of the newly approved agents. Obesity, being a complex multifactorial disease with several pathways disturbed, would require a molecule or a combination of molecules to act at multiple pathways/targets if it is to produce clinically significant weight reduction. There are several instances of older drugs approved for other indications being explored such as zonisamide, naltrexone, and bupropion. Some of the new targets that have evoked great interest in the last few years include SIRT1, GLP-1, aminopeptidase-2, sodium glucose cotransporter, and NPY, but much needs to be done before effective drugs are discovered that target these receptors. It can be safely surmised that as our knowledge of the pathophysiology of obesity improves, it may be possible to discover the “holy grail” of antiobesity drugs.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

1. WHO. Obesity and Overweight [Internet]; 2013. <http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed May 23, 2013.
2. Wolf AM, Colditz GA. Current estimates of the economic cost of obesity in the United States. *Obes Res*. 1998;6(2):97-106.
3. Kumanyika SK, Obarzanek E, Stettler N, et al Population-based prevention of obesity: the need for comprehensive promotion of healthful eating, physical activity, and energy balance: a scientific statement from American heart association council on epidemiology and prevention, interdisciplinary committee for prevention (formerly the expert panel on population and prevention science). *Circulation*. 2008;118(4):428-464.
4. Obesity to Overtake Smoking as Leading Cause of Death [Internet]; 2013. <http://www.medscape.com/viewarticle/474966>. Accessed May 31, 2013.
5. Lavie CJ, Milani RV, Ventura HO. Obesity and cardiovascular disease: risk factor, paradox, and impact of weight loss. *J Am Coll Cardiol*. 2009;53(21):1925-1932.
6. Kremers S, Reubsat A, Martens M, et al Systematic prevention of overweight and obesity in adults: a qualitative and quantitative literature analysis. *Obes Rev*. 2010;11(5):371-379.
7. Flum DR, Salem L, Elrod JAB, Dellinger EP, Cheadle A, Chan L. Early mortality among medicare beneficiaries undergoing bariatric surgical procedures. *JAMA*. 2005;294(15):1903-1908.
8. Wolfe BM, Morton JM. Weighing in on bariatric surgery: procedure use, readmission rates, and mortality. *JAMA*. 2005;294(15):1960-1963.
9. Filippatos TD, Derdemezis CS, Gazi IF, Nakou ES, Mikhailidis DP, Elisaf MS. Orlistat-associated adverse effects and drug interactions: a critical review. *Drug Saf*. 2008;31(1):53-65.
10. McKay RH. Long-term use of diethylpropion in obesity. *Curr Med Res Opin*. 1973;1(8):489-493.
11. Ioannides-Demos LL, Proietto J, Tonkin AM, McNeil JJ. Safety of drug therapies used for weight loss and treatment of obesity. *Drug Saf*. 2006;29(4):277-302.
12. Bray GA, Greenway FL. Pharmacological treatment of the overweight patient. *Pharmacol Rev*. 2007;59(2):151-184.
13. BELVIQ—022529lbl.pdf [Internet]; 2013. [http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2012/022529lbl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/022529lbl.pdf). Accessed May 23, 2013.
14. Thomsen WJ, Grottick AJ, Menzaghi F, et al Lorcaserin, a novel selective human 5-hydroxytryptamine 2C agonist: in vitro and in vivo pharmacological characterization. *J Pharmacol Exp Ther*. 2008;325(2):577-587.
15. Weissman NJ, Tighe JF, Jr, Gottdiener JS, Gwynne JT. An assessment of heart-valve abnormalities in obese patients taking dexfenfluramine, sustained-release dexfenfluramine, or placebo. sustained-release dexfenfluramine study group. *N Engl J Med*. 1998;339(11):725-732.
16. Sadeque AJM, Usmani KA, Palamar S, Cerny MA, Chen WG. Identification of human UDP-glucuronosyltransferases involved in N-carbamoyl glucuronidation of lorcaserin. *Drug Metab Dispos*. 2012;40(4):772-778.
17. Usmani KA, Chen WG, Sadeque AJM. Identification of human cytochrome P450 and flavin-containing monooxygenase enzymes involved in the metabolism of lorcaserin, a novel selective human 5-hydroxytryptamine 2C agonist. *Drug Metab Dispos*. 2012;40(4):761-771.
18. Smith SR, Weissman NJ, Anderson CM, et al Multicenter, placebo-controlled trial of lorcaserin for weight management. *N Engl J Med*. 2010;363(3):245-256.
19. Fidler MC, Sanchez M, Raether B, et al A one-year randomized trial of lorcaserin for weight loss in obese and overweight adults: the BLOSSOM trial. *J Clin Endocrinol Metab*. 2011;96(10):3067-3077.
20. O'Neil PM, Smith SR, Weissman NJ, et al Randomized placebo-controlled clinical trial of lorcaserin for weight loss in type 2

- diabetes mellitus: the BLOOM-DM study. *Obesity (Silver Spring)*. 2012;20(7):1426-1436.
21. Persistent Concerns About Lorcaserin (Lorqess) From FDA Reviewers [Internet]. Forbes; 2013. <http://www.forbes.com/sites/larryhusten/2012/05/08/persistent-concerns-about-lorcaserin-lorqess-from-fda-reviewers/>. Accessed May 24, 2013.
  22. Press Announcements > FDA approves Belviq to treat some overweight or obese adults [Internet]; 2013. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm309993.htm>. Accessed May 26, 2013.
  23. Allison DB, Gadde KM, Garvey WT, et al Controlled-release phentermine/topiramate in severely obese adults: a randomized controlled trial (EQUIP). *Obesity (Silver Spring)*. 2012;20(2):330-342.
  24. Gadde KM, Allison DB, Ryan DH, et al Effects of low-dose, controlled-release, phentermine plus topiramate combination on weight and associated comorbidities in overweight and obese adults (CONQUER): a randomised, placebo-controlled, phase 3 trial. *Lancet*. 2011;377(9774):1341-1352.
  25. Garvey WT, Ryan DH, Look M, et al Two-year sustained weight loss and metabolic benefits with controlled-release phentermine/topiramate in obese and overweight adults (SEQUEL): a randomized, placebo-controlled, phase 3 extension study. *Am J Clin Nutr*. 2012;95(2):297-308.
  26. Uwaifo G, Cosentino, Conrad A. Phentermine and topiramate for the management of obesity: a review. *Drug Des Devel Ther*. 2011; 7:267-278.
  27. Risk Evaluation and Mitigation Strategy (REMS). Qsymia® (phentermine and topiramate extended-release) capsules CIV [Internet]; 2013. <http://www.qsymiarems.com/>. Accessed May 31, 2013.
  28. Wadden TA, Foreyt JP, Foster GD, et al Weight loss with naltrexone SR/bupropion SR combination therapy as an adjunct to behavior modification: the COR-BMOD trial. *Obesity (Silver Spring)*. 2011;19(1):110-120.
  29. Greenway FL, Fujioka K, Plodkowski RA, et al Effect of naltrexone plus bupropion on weight loss in overweight and obese adults (COR-I): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2010;376(9741):595-605.
  30. Apovian C, Aronne L, Rubino D, et al A randomized, phase 3 trial of naltrexone SR/bupropion SR on weight and obesity-related risk factors (COR-II). *Obesity (Silver Spring)*. 2013;21(5):935-943.
  31. Orexigen Therapeutics, Inc.—Contrave [Internet]; 2013. <http://www.orexigen.com/product-candidates/contrave.html>. Accessed June 3, 2013.
  32. Gadde KM, Yonish GM, Foust MS, Wagner HR. Combination therapy of zonisamide and bupropion for weight reduction in obese women: a preliminary, randomized, open-label study. *J Clin Psychiatry*. 2007;68(8):1226-1229.
  33. Powell AG, Apovian CM, Aronne LJ. New drug targets for the treatment of obesity. *Clin Pharmacol Ther*. 2011;90(1):40-51.
  34. Orexigen Therapeutics, Inc.—Press Release [Internet]; 2013. <http://ir.orexigen.com/phoenix.zhtml?c=207034&p=irol-newsArticle&id=1400316>. Accessed May 23, 2013.
  35. Nielsen A-LH, Larsen TM, Madsbad S, et al The effect of tesofensine on body weight and body composition in obese subjects—secondary publication [in Danish]. *Ugeskr Laeger*. 2009;171(41):2974-2977.
  36. Doggrel SA. Tesofensine—a novel potent weight loss medicine. Evaluation of: Astrup A, Breum L, Jensen TJ, Kroustrup JP, Larsen TM. Effect of tesofensine on bodyweight loss, body composition, and quality of life in obese patients: a randomised, double-blind, placebo-controlled trial. *Lancet* 2008;372:1906-13. *Expert Opin Investig Drugs*. 2009;18(7):1043-1046.
  37. Trial irregularities earn Lancet study of potential weight loss drug tesofensine Expression of Concern. Retraction Watch [Internet]; 2013. <http://retractionwatch.wordpress.com/2013/04/09/trial-irregularities-earn-lancet-study-of-potential-weight-loss-drug-tesofensine-expression-of-concern/>. Accessed May 23, 2013.
  38. Yeomans MR, Wright P. Lower pleasantness of palatable foods in nalmefene-treated human volunteers. *Appetite*. 1991;16(3):249-259.
  39. Yeomans MR, Gray RW. Opioid peptides and the control of human ingestive behaviour. *Neurosci Biobehav Rev*. 2002;26(6):713-728.
  40. Nathan PJ, O'Neill BV, Napolitano A, Bullmore ET. Neuropsychiatric adverse effects of centrally acting antiobesity drugs. *CNS Neurosci Ther*. 2011;17(5):490-505.
  41. Nathan PJ, Bush MA, Tao WX, et al Multiple-dose safety, pharmacokinetics, and pharmacodynamics of the  $\mu$ -opioid receptor inverse agonist GSK1521498. *J Clin Pharmacol*. 2012;52(10):1456-1467.
  42. Kopelman P, Bryson A, Hickling R, et al Cetilistat (ATL-962), a novel lipase inhibitor: a 12-week randomized, placebo-controlled study of weight reduction in obese patients. *Int J Obes (Lond)*. 2007;31(3):494-499.
  43. Kopelman P, Groot G de H, Rissanen A, et al Weight loss, HbA1c reduction, and tolerability of cetilistat in a randomized, placebo-controlled phase 2 trial in obese diabetics: comparison with orlistat (Xenical). *Obesity (Silver Spring)*. 2010;18(1):108-115.
  44. Valentino MA, Lin JE, Waldman SA. Central and peripheral molecular targets for antiobesity pharmacotherapy. *Clin Pharmacol Ther*. 2010;87(6):652-662.
  45. Astrup A, Rössner S, Van Gaal L, et al Effects of liraglutide in the treatment of obesity: a randomised, double-blind, placebo-controlled study. *Lancet*. 2009;374(9701):1606-1616.
  46. Search of: liraglutide AND obesity—List Results—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/results?term=liraglutide+AND+obesity&Search=Search>. Accessed May 23, 2013.
  47. Vilsbøll T, Christensen M, Junker AE, Knop FK, Gluud LL. Effects of glucagon-like peptide-1 receptor agonists on weight loss: systematic review and meta-analyses of randomised controlled trials. *BMJ*. 2012;344:d7771.
  48. Maciejewski BS, Laperle JL, Chen D, et al Pharmacological inhibition to examine the role of DGAT1 in dietary lipid absorption in rodents and humans. *Am J Physiol Gastrointest Liver Physiol*. 2013;304(11):G958-G969.
  49. Search of: PF-04620110—List Results—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/results?term=PF-04620110&Search=Search>. Accessed May 28, 2013.
  50. The Science—Our Approach—Zafgen [Internet]; 2013. <http://zafgen.com/zafgen/our-approach/the-science>. Accessed May 24, 2013.
  51. Hughes TE, Kim DD, Marjason J, Proietto J, Whitehead JP, Vath JE. Ascending dose controlled trial of beloranib, a novel obesity

- treatment for safety, tolerability and weight loss in obese women [published online March 20, 2013]. *Obes (Silver Spring)*. 2013.
52. Search of: beloranib—List Results—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/results?term=beloranib&Search=Search>. Accessed May 24, 2013.
  53. Timmers S, Konings E, Bilet L, et al Calorie restriction-like effects of 30 days of resveratrol supplementation on energy metabolism and metabolic profile in obese humans. *Cell Metab*. 2011; 14(5):612-622.
  54. Brakenhielm E, Cao Y. Angiogenesis in adipose tissue. *Methods Mol Biol*. 2008;456:65-81.
  55. Christiaens V, Lijnen HR. Angiogenesis and development of adipose tissue. *Mol Cell Endocrinol*. 2010;318(1-2):2-9.
  56. Lijnen HR. Angiogenesis and obesity. *Cardiovasc Res*. 2008; 78(2):286-293.
  57. Silha JV, Krsek M, Sucharda P, Murphy LJ. Angiogenic factors are elevated in overweight and obese individuals. *Int J Obes (Lond)*. 2005;29(11):1308-1314.
  58. npre20116256-1.pdf [Internet]; 2013. <http://precedings.nature.com/documents/6256/version/1/files/npre20116256-1.pdf>. Accessed May 28, 2013.
  59. nyhf2\_10\_hottechnologies\_angiolab.pdf [Internet]; 2013. [http://www.nyhf.org/docs/nyhf2\\_10\\_hottechnologies\\_angiolab.pdf](http://www.nyhf.org/docs/nyhf2_10_hottechnologies_angiolab.pdf). Accessed May 28, 2013.
  60. Yoon M, Kim M-Y. The anti-angiogenic herbal composition Ob-X from *Morus alba*, *Melissa officinalis*, and *Artemisia capillaris* regulates obesity in genetically obese ob/ob mice. *Pharm Biol*. 2011;49(6):614-619.
  61. Takeda Submits a New Drug Application for Cetilistat (Development Code: ATL-962) in Japan for the Treatment of Obesity with Complications. Takeda Pharmaceutical Company Limited [Internet]; 2013. [http://www.takeda.com/news/2012/20121030\\_4004.html](http://www.takeda.com/news/2012/20121030_4004.html). Accessed June 1, 2013.
  62. Cardiovascular Outcomes Study of Naltrexone SR/Bupropion SR in Overweight and Obese Subjects With Cardiovascular Risk Factors (The Light Study)—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://clinicaltrials.gov/ct2/show/NCT01601704?term=contrave&rank=3>. Accessed June 1, 2013.
  63. Orexigen Therapeutics, Inc.—Empatic [Internet]; 2013. <http://www.orexigen.com/product-candidates/empatic.html>. Accessed June 1, 2013.
  64. Effect of Liraglutide on Body Weight in Non-diabetic Obese Subjects or Overweight Subjects With Co-morbidities: SCALE™—Obesity and Pre-diabetes—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://clinicaltrials.gov/ct2/show/NCT01272219?term=Liraglutide+AND+obesity&rank=1>. Accessed June 1, 2013.
  65. Search of: velneperit—List Results—ClinicalTrials.gov [Internet]; 2013. <http://clinicaltrials.gov/ct2/results?term=velneperit&Search=Search>. Accessed June 1, 2013.
  66. Search of: tesofensine—List Results—ClinicalTrials.gov [Internet]; 2013. [http://www.clinicaltrials.gov/ct2/results/displayOpt?flds=a&flds=b&flds=f&submit\\_fld\\_opt=on&term=tesofensine&show\\_flds=Y](http://www.clinicaltrials.gov/ct2/results/displayOpt?flds=a&flds=b&flds=f&submit_fld_opt=on&term=tesofensine&show_flds=Y). Accessed June 1, 2013.
  67. A Study of the Safety and Effectiveness of JNJ-16269110 (R256918) in Overweight and Obese Patients—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://clinicaltrials.gov/ct2/show/NCT00622765>. Accessed June 1, 2013.
  68. Safety Study of the Inhibition of Agouti-related Protein (AgRP) for the Management of Obesity and Weight Loss—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/show/NCT00779519?term=TTP435&rank=1>. Accessed June 3, 2013.
  69. Wynne K, Park AJ, Small CJ, et al Subcutaneous oxyntomodulin reduces body weight in overweight and obese subjects: a double-blind, randomized, controlled trial. *Diabetes*. 2005;54(8): 2390-2395.
  70. Search of: GSK 1521498—List Results—ClinicalTrials.gov [Internet]; 2013. [http://www.clinicaltrials.gov/ct2/results/displayOpt?flds=a&flds=b&flds=f&submit\\_fld\\_opt=on&term=GSK+1521498&show\\_flds=Y](http://www.clinicaltrials.gov/ct2/results/displayOpt?flds=a&flds=b&flds=f&submit_fld_opt=on&term=GSK+1521498&show_flds=Y). Accessed May 27, 2013.
  71. Effects of the D3 Antagonist GSK598809 on Food Reward and Reinforcement—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/show/NCT01039454?term=GSK+598809+AND+obesity&rank=1>. Accessed June 1, 2013.
  72. Search of: PF-04971729 AND obesity—List Results—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/results?term=PF-04971729+AND+obesity&Search=Search>. Accessed June 1, 2013.
  73. Search of: PF-04620110 AND obesity—List Results—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/results?term=PF-04620110+AND+obesity&Search=Search>. Accessed June 1, 2013.
  74. Cooke D, Bloom S. The obesity pipeline: current strategies in the development of anti-obesity drugs. *Nat Rev Drug Discov*. 2006; 5(11):919-931.
  75. Greenway FL, Caruso MK. Safety of obesity drugs. *Expert Opin Drug Saf*. 2005;4(6):1083-1095.
  76. Ioannides-Demos LL, Proietto J, McNeil JJ. Pharmacotherapy for obesity. *Drugs*. 2005;65(10):1391-1418.
  77. Abenhaim L, Moride Y, Brenot F, et al. Appetite-suppressant drugs and the risk of primary pulmonary hypertension. International primary pulmonary hypertension study group. *N Engl J Med*. 1996;335(9):609-616.
  78. Jick H, Vasilakis C, Weinrauch LA, Meier CR, Jick SS, Derby LE. A population-based study of appetite-suppressant drugs and the risk of cardiac-valve regurgitation. *N Engl J Med* 1998; 339(11):719-724.
  79. Sachdev M, Miller WC, Ryan T, Jollis JG. Effect of fenfluramine-derivative diet pills on cardiac valves: a meta-analysis of observational studies. *Am Heart J*. 2002;144(6):1065-1073.
  80. Glazer G. Long-term pharmacotherapy of obesity 2000: a review of efficacy and safety. *Arch Intern Med*. 2001;161(15):1814-1824.
  81. Kernan WN, Viscoli CM, Brass LM, et al Phenylpropranolamine and the risk of hemorrhagic stroke. *N Engl J Med*. 2000;343(25): 1826-1832.
  82. Li Z, Maglione M, Tu W, et al Meta-analysis: pharmacologic treatment of obesity. *Ann Intern Med*. 2005;142(7):532-546.
  83. Uretsky S, Messerli FH, Bangalore S, et al Obesity paradox in patients with hypertension and coronary artery disease. *Am J Med*. 2007;120(10):863-870.

84. Wassertheil-Smoller S, Fann C, Allman RM, et al Relation of low body mass to death and stroke in the systolic hypertension in the elderly program. The SHEP cooperative research group. *Arch Intern Med*. 2000;160(4):494-500.
85. Lavie CJ, Alpert MA, Arena R, Mehra MR, Milani RV, Ventura HO. Impact of obesity and the obesity paradox on prevalence and prognosis in heart failure. *JACC Heart Fail*. 2013;1(2):93-102.
86. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA*. 2013;309(1):71-82.
87. Fonarow GC, Srikanthan P, Costanzo MR, Cintron GB, Lopatin M, ADHERE Scientific Advisory Committee and Investigators. An obesity paradox in acute heart failure: analysis of body mass index and inhospital mortality for 108,927 patients in the acute decompensated heart failure national registry. *Am Heart J*. 2007;153(1):74-81.
88. Artham SM, Lavie CJ, Milani RV, Ventura HO. Value of weight reduction in patients with cardiovascular disease. *Curr Treat Options Cardiovasc Med*. 2010;12(1):21-35.
89. Christensen R, Kristensen PK, Bartels EM, Bliddal H, Astrup A. Efficacy and safety of the weight-loss drug rimonabant: a meta-analysis of randomised trials. *Lancet*. 2007;370(9600):1706-1713.
90. Mitchell PB, Morris MJ. Depression and anxiety with rimonabant. *Lancet*. 2007;370(9600):1671-1672.
91. James WPT, Caterson ID, Coutinho W, et al Effect of sibutramine on cardiovascular outcomes in overweight and obese subjects. *N Engl J Med*. 2010;363(10):905-917.
92. Amylin and Takeda Discontinue Development of Pramlintide/Metreleptin Combination Treatment for Obesity Following Commercial Reassessment of the Program. Takeda Pharmaceutical Company Limited [Internet]; 2013. [http://www.takeda.com/news/2011/20110805\\_3889.html](http://www.takeda.com/news/2011/20110805_3889.html). Accessed June 3, 2013.
93. Proefschrift Edwin Parlevliet 180810-06.pdf [Internet]; 2013. <https://openaccess.leidenuniv.nl/bitstream/handle/1887/16080/06.pdf?sequence=9>. Accessed May 24, 2013.
94. A Weight Loss Study in Overweight Men and Women—Full Text View—ClinicalTrials.gov [Internet]; 2013. <http://www.clinicaltrials.gov/ct2/show/NCT00993421?term=LY377604&rank=1>. Accessed June 3, 2013.
95. Barak N, Greenway FL, Fujioka K, Aronne LJ, Kushner RF. Effect of histaminergic manipulation on weight in obese adults: a randomized placebo controlled trial. *Int J Obes (Lond)*. 2008;32(10):1559-1565.
96. Ali AH, Yanoff LB, Stern EA, et al Acute effects of betahistine hydrochloride on food intake and appetite in obese women: a randomized, placebo-controlled trial. *Am J Clin Nutr*. 2010;92(6):1290-1297.
97. Aronne LJ, Tonstad S, Moreno M, et al. A clinical trial assessing the safety and efficacy of taranabant, a CB1R inverse agonist, in obese and overweight patients: a high-dose study. *Int J Obes (Lond)*. 2010;34(5):919-935.
98. Krishna R, Gumbiner B, Stevens C, et al Potent and selective agonism of the melanocortin receptor 4 with MK-0493 does not induce weight loss in obese human subjects: energy intake predicts lack of weight loss efficacy. *Clin Pharmacol Ther*. 2009;86(6):659-66.
99. Heal DJ, Gosden J, Smith SL. Regulatory challenges for new drugs to treat obesity and comorbid metabolic disorders. *Br J Clin Pharmacol*. 2009;68(6):861-874.
100. Guideline on Weight Control—Adopted—WC500003264.pdf [Internet]; 2013. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC500003264.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003264.pdf). Accessed June 3, 2013.
101. Guidance for Industry Developing Products for Weight Management—ucm071612.pdf [Internet]; 2013. <http://www.fda.gov/downloads/Drugs/.../Guidances/ucm071612.pdf>. Accessed June 3, 2013.